

INSTRUCTIONS FOR USE

इस्तेमाल की जानकारी



Natjewel FS® Thiamethoxam 30% FS Systemic insecticide

नॅटजेवेल एफएस थायामेथोक्सम 30% एफ एस सिस्टमिक एकाइस

नॅटजेवेल एफएस थायामेथोक्सम 30% एफ एस सिस्टमिक एकाइस

FOR AGRICULTURAL USE ONLY.

Manufactured and Marketed by: NATCO Pharms. Pvt. Ltd., Bangalore Hills, Hyderabad - 500 024, Telangana, India.


उत्पादक और बाजार में लाया गया: NATCO फार्मास प्रिवेट लिमिटेड, बंगलूर हिल्स, हैदराबाद - 500 024, तेलंगणा, भारत।

Customer Care: Tel: 040 23545561 / 23600836. Email: crophealth@natcoapharma.com

KEEP IN COOL, DRY PLACE AWAY FROM HEAT AND OPEN FLAME

نیت جیول ایف ایس

ثیامیٹوکسام 30% ایف ایس



Natjewel FS® Thiamethoxam 30% FS Systemic insecticide

نॅٹजेوئل ایف ایس

ثیامیٹوکسام 30% ایف ایس

FOR AGRICULTURAL USE ONLY.

Manufactured and Marketed by: NATCO Pharms. Pvt. Ltd., Bangalore Hills, Hyderabad - 500 024, Telangana, India.


उत्पादक और बाजार में लाया गया: NATCO फार्मास प्रिवेट लिमिटेड, बंगलूर हिल्स, हैदराबाद - 500 024, तेलंगणा, भारत।

Customer Care: Tel: 040 23545561 / 23600836. Email: crophealth@natcoapharma.com

KEEP IN COOL, DRY PLACE AWAY FROM HEAT AND OPEN FLAME

നാട്ടുജവേൽ ഏഫ്എസ്

ത്യാഥിമെതോക്സാം 30% ഏഫ്.എസ്.



Natjewel FS® Thiamethoxam 30% FS Systemic insecticide

നാട്ടുജവേൽ എഫ്എസ്

ത്യാഥിമെതോക്സാം 30% ഏഫ്.എസ്.

FOR AGRICULTURAL USE ONLY.

Manufactured and Marketed by: NATCO Pharms. Pvt. Ltd., Bangalore Hills, Hyderabad - 500 024, Telangana, India.

उत्पादक और बाजार में लाया गया: NATCO फार्मास प्रिवेट लिमिटेड, बंगलूर हिल्स, हैदराबाद - 500 024, तेलंगणा, भारत।

Customer Care: Tel: 040 23545561 / 23600836. Email: crophealth@natcoapharma.com

KEEP IN COOL, DRY PLACE AWAY FROM HEAT AND OPEN FLAME

நாட்டுஜவீல் ஏஃபிஎஃஸ்

த்யாதிமெதொக்ஸாம் 30% ஏஃபி.எஃஸ்.



Natjewel FS® Thiamethoxam 30% FS Systemic insecticide

நாட்டுஜவீல் ஏஃபிஎஃஸ்

த்யாதிமெதொக்ஸாம் 30% ஏஃபி.எஃஸ்.

FOR AGRICULTURAL USE ONLY.

Manufactured and Marketed by: NATCO Pharms. Pvt. Ltd., Bangalore Hills, Hyderabad - 500 024, Telangana, India.


उत्पादक और बाजार में लाया गया: NATCO फार्मास प्रिवेट लिमिटेड, बंगलूर हिल्स, हैदराबाद - 500 024, तेलंगणा, भारत।

Customer Care: Tel: 040 23545561 / 23600836. Email: crophealth@natcoapharma.com

KEEP IN COOL, DRY PLACE AWAY FROM HEAT AND OPEN FLAME

நட்டுஜவீல் ஏஃபிஎஃஸ்

த்யாதிமெதொக்ஸாம் 30% ஃபி.எஃஸ்.



Natjewel FS® Thiamethoxam 30% FS Systemic insecticide

நட்டுஜவீல் ஏஃபிஎஃஸ்

த்யாதிமெதொக்ஸாம் 30% ஃபி.எஃஸ்.

FOR AGRICULTURAL USE ONLY.

Manufactured and Marketed by: NATCO Pharms. Pvt. Ltd., Bangalore Hills, Hyderabad - 500 024, Telangana, India.

उत्पादक और बाजार में लाया गया: NATCO फार्मास प्रिवेट लिमिटेड, बंगलूर हिल्स, हैदराबाद - 500 024, तेलंगणा, भारत।

Customer Care: Tel: 040 23545561 / 23600836. Email: crophealth@natcoapharma.com

KEEP IN COOL, DRY PLACE AWAY FROM HEAT AND OPEN FLAME

Natjewel FS® Thiamethoxam 30% FS Systemic Insecticide

Thiamethoxam 30% FS is a broad-spectrum systemic insecticide that is well suited for seed treatment to control early season sucking pests...

Table with 4 columns: Crops, Common Name of Pest, Dosage (AI per kg seed), and Waiting period before last spray to harvest (days). Rows include Cotton, Sorghum, Rice, Maize, etc.

Directions of Use: For all crops, use the required quantity of Thiamethoxam 30% FS with 10-20 ml water for each kg of seed to be treated...

Application Procedure: 1. Prepare this Slurry as described. 2. Fill the seed into the receptacle. 3. Start mixing the seed. 4. Apply the required amount of slurry pouring into the mixing seed...

Precaution: Do not use cooling utensils for preparing slurry. Avoid spreading the slurry solution. Avoid contact with skin, eyes and clothing...

Symptoms of Poisoning: Possessing may cause symptoms of dizziness, faintness, nausea, vomiting, diarrhoea, convulsions and fits.

First Aid: Remove the affected person to a well-ventilated area or fresh air and protect him from overexposure. In case of Skin Contact: Remove contaminated clothing and thoroughly wash the affected parts of the body with soap and water...

Disposal: Use non-hazardous incineration. Do not burn in open air. Do not dump in water bodies. Do not dump in landfills.

Storage Conditions: The packages containing the insecticides should be stored in original containers in a cool, dry place...

Table with 2 columns: Insecticide Name and Concentration. Rows include Chlorantraniliprole, Thiamethoxam, Imidacloprid, etc.

Warning: 1. Do not use on crops other than those specified on the label. 2. Destroy the container after use as directed on label.

നാട്ടുജവേൽ എഫ്എസ്

ത്യാഥിമെതോക്സാം 30% ഏഫ്.എസ്. (സിസ്റ്റമിക് ഇൻസെക്റ്റിസൈഡ്)

Table with 4 columns: Crops, Common Name of Pest, Dosage (AI per kg seed), and Waiting period before last spray to harvest (days). Rows include Cotton, Sorghum, Rice, Maize, etc.

Directions of Use: For all crops, use the required quantity of Thiamethoxam 30% FS with 10-20 ml water for each kg of seed to be treated...

Application Procedure: 1. Prepare this Slurry as described. 2. Fill the seed into the receptacle. 3. Start mixing the seed. 4. Apply the required amount of slurry pouring into the mixing seed...

Precaution: Do not use cooling utensils for preparing slurry. Avoid spreading the slurry solution. Avoid contact with skin, eyes and clothing...

Symptoms of Poisoning: Possessing may cause symptoms of dizziness, faintness, nausea, vomiting, diarrhoea, convulsions and fits.

First Aid: Remove the affected person to a well-ventilated area or fresh air and protect him from overexposure. In case of Skin Contact: Remove contaminated clothing and thoroughly wash the affected parts of the body with soap and water...

Disposal: Use non-hazardous incineration. Do not burn in open air. Do not dump in water bodies. Do not dump in landfills.

Storage Conditions: The packages containing the insecticides should be stored in original containers in a cool, dry place...

Table with 2 columns: Insecticide Name and Concentration. Rows include Chlorantraniliprole, Thiamethoxam, Imidacloprid, etc.

Warning: 1. Do not use on crops other than those specified on the label. 2. Destroy the container after use as directed on label.

نٹ جیول ایف ایس

ثیامیٹوکسام 30% ایف ایس (سیسٹمیک اینٹی سیکٹو)

Table with 4 columns: Crops, Common Name of Pest, Dosage (AI per kg seed), and Waiting period before last spray to harvest (days). Rows include Cotton, Sorghum, Rice, Maize, etc.

Directions of Use: For all crops, use the required quantity of Thiamethoxam 30% FS with 10-20 ml water for each kg of seed to be treated...

Application Procedure: 1. Prepare this Slurry as described. 2. Fill the seed into the receptacle. 3. Start mixing the seed. 4. Apply the required amount of slurry pouring into the mixing seed...

Precaution: Do not use cooling utensils for preparing slurry. Avoid spreading the slurry solution. Avoid contact with skin, eyes and clothing...

Symptoms of Poisoning: Possessing may cause symptoms of dizziness, faintness, nausea, vomiting, diarrhoea, convulsions and fits.

First Aid: Remove the affected person to a well-ventilated area or fresh air and protect him from overexposure. In case of Skin Contact: Remove contaminated clothing and thoroughly wash the affected parts of the body with soap and water...

Disposal: Use non-hazardous incineration. Do not burn in open air. Do not dump in water bodies. Do not dump in landfills.

Storage Conditions: The packages containing the insecticides should be stored in original containers in a cool, dry place...

Table with 2 columns: Insecticide Name and Concentration. Rows include Chlorantraniliprole, Thiamethoxam, Imidacloprid, etc.

Warning: 1. Do not use on crops other than those specified on the label. 2. Destroy the container after use as directed on label.

नॅटजेवेल एफएस

थायामेथोक्सम 30% एफ एस (सिस्टमिक इन्सेक्टिसाइड)

Table with 4 columns: Crops, Common Name of Pest, Dosage (AI per kg seed), and Waiting period before last spray to harvest (days). Rows include Cotton, Sorghum, Rice, Maize, etc.

Directions of Use: For all crops, use the required quantity of Thiamethoxam 30% FS with 10-20 ml water for each kg of seed to be treated...

Application Procedure: 1. Prepare this Slurry as described. 2. Fill the seed into the receptacle. 3. Start mixing the seed. 4. Apply the required amount of slurry pouring into the mixing seed...

Precaution: Do not use cooling utensils for preparing slurry. Avoid spreading the slurry solution. Avoid contact with skin, eyes and clothing...

Symptoms of Poisoning: Possessing may cause symptoms of dizziness, faintness, nausea, vomiting, diarrhoea, convulsions and fits.

First Aid: Remove the affected person to a well-ventilated area or fresh air and protect him from overexposure. In case of Skin Contact: Remove contaminated clothing and thoroughly wash the affected parts of the body with soap and water...

Disposal: Use non-hazardous incineration. Do not burn in open air. Do not dump in water bodies. Do not dump in landfills.

Storage Conditions: The packages containing the insecticides should be stored in original containers in a cool, dry place...

Table with 2 columns: Insecticide Name and Concentration. Rows include Chlorantraniliprole, Thiamethoxam, Imidacloprid, etc.

Warning: 1. Do not use on crops other than those specified on the label. 2. Destroy the container after use as directed on label.

नॅटजेवेल एफएस

थायामेथोक्सम 30% एफ एस (सिस्टमिक इन्सेक्टिसाइड)

Table with 4 columns: Crops, Common Name of Pest, Dosage (AI per kg seed), and Waiting period before last spray to harvest (days). Rows include Cotton, Sorghum, Rice, Maize, etc.

Directions of Use: For all crops, use the required quantity of Thiamethoxam 30% FS with 10-20 ml water for each kg of seed to be treated...

Application Procedure: 1. Prepare this Slurry as described. 2. Fill the seed into the receptacle. 3. Start mixing the seed. 4. Apply the required amount of slurry pouring into the mixing seed...

Precaution: Do not use cooling utensils for preparing slurry. Avoid spreading the slurry solution. Avoid contact with skin, eyes and clothing...

Symptoms of Poisoning: Possessing may cause symptoms of dizziness, faintness, nausea, vomiting, diarrhoea, convulsions and fits.

First Aid: Remove the affected person to a well-ventilated area or fresh air and protect him from overexposure. In case of Skin Contact: Remove contaminated clothing and thoroughly wash the affected parts of the body with soap and water...

Disposal: Use non-hazardous incineration. Do not burn in open air. Do not dump in water bodies. Do not dump in landfills.

Storage Conditions: The packages containing the insecticides should be stored in original containers in a cool, dry place...

Table with 2 columns: Insecticide Name and Concentration. Rows include Chlorantraniliprole, Thiamethoxam, Imidacloprid, etc.

Warning: 1. Do not use on crops other than those specified on the label. 2. Destroy the container after use as directed on label.

1852 Natjewel FS® DFU Size: 329.0 mm (W) x 180.91 mm (H)